

SYMMETRY BREAKING FOR SCHRÖDINGER-POISSON-SLATER ENERGY

JACOPO BELLAZZINI AND MARCO GHIMENTI

ABSTRACT. We study the asymptotic behavior of ground state energy for Schrödinger-Poisson-Slater energy functional. We show that ground state energy restricted to radially symmetric functions is above the ground state energy when the number of particles is sufficiently large.

The aim of this paper is to show symmetry breaking phenomena for the ground states energy of the following Schrödinger-Poisson-Slater (SPS) equation,

$$(0.1) \quad i\psi_t + \Delta\psi - (|x|^{-1} * |\psi|^2)\psi + |\psi|^{\frac{2}{3}}\psi = 0 \quad \text{in } \mathbb{R}^3,$$

where $\psi(x, t) : \mathbb{R}^3 \times [0, T) \rightarrow \mathbb{C}$ is the wave function, $*$ denotes the convolution, and the nonlinear term $|\psi|^{\frac{2}{3}}\psi$ is the Slater correction term.

This Schrödinger-type equation with a repulsive nonlocal Coulomb potential is obtained by approximation of the Hartree-Fock equation describing a quantum mechanical system of many particles, see e.g. [1, 12, 7]. As example, SPS equation gives a good approximation of the time evolution of an electron ensemble in a semiconductor crystal. In particular, in this mean field approximation the quantity $\int |\psi|^2 dx = \rho^2$ describes the total amount of particles. The energy associated to a state ψ is defined as

$$E(\psi) := \frac{1}{2} \|\nabla\psi\|_{L^2}^2 + \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|\psi(x)|^2 |\psi(y)|^2}{|x - y|} dx dy - \frac{3}{8} \int_{\mathbb{R}^3} |\psi|^{\frac{8}{3}} dx.$$

We define the *ground state energy* the quantity I_{ρ^2} , defined as

$$I_{\rho^2} = \inf_{B_{\rho}} E(\psi)$$

where $B_{\rho} := \{\psi \in H^1(\mathbb{R}^3) \text{ such that } \|\psi\|_2^2 = \rho^2\}$.

Very briefly we summarize what is known about I_{ρ^2} (see e.g [6], [5], [4], [10], [15], [8])

- (i) $-\infty < I_{\rho^2} < 0$ for all $\rho > 0$
- (ii) the ground state energy is weakly subadditive, namely that

$$I_{\rho^2} \leq I_{\mu^2} + I_{\rho^2 - \mu^2} \text{ for all } 0 < \mu < \rho$$

- (iii) the function $\rho \rightarrow I_{\rho^2}$ is continuous

Key words and phrases. Symmetry breaking, Schrödinger-Poisson-Slater energy, constrained minimization.

- (iv) $\lim_{\rho \rightarrow 0} \frac{I_{\rho^2}}{\rho^2} = 0$
 - (v) minimizers for E on B_ρ exist when ρ is sufficiently small
 - (vi) minimizers for E on B_ρ are radially symmetric when ρ is sufficiently small
- If we define

$$I_{\rho^2}^{rad} = \inf_{B_\rho^{rad}} E(\psi)$$

where $B_\rho^{rad} := \{\psi \in H^1(\mathbb{R}^3) \text{ radially symmetric such that } \|\psi\|_2^2 = \rho^2\}$ it is not clear if

$$I_{\rho^2}^{rad} = I_{\rho^2} \text{ for all } \rho > 0. \quad (Q)$$

We shall emphasize that for Schrödinger-Poisson-Slater energy there are two terms that are in competition when we move from u to $u^\star$, the symmetric decreasing rearrangement of u . Indeed, it is well understood that for kinetic term

$$\int_{\mathbb{R}^3} |\nabla u^\star|^2 dx \leq \int_{\mathbb{R}^3} |\nabla u|^2 dx$$

while for the Coulomb term thanks to Riesz inequality

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|u(x)|^2 |u(y)|^2}{|x-y|} dx dy \leq \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|u^\star(x)|^2 |u^\star(y)|^2}{|x-y|} dx dy$$

such that the question (Q) is still open.

Aim of this paper is to give an answer to question (Q). The fact that the energy functional as well as the L^2 constraint is rotationally invariant does not imply that the ground state energy for radially symmetric functions coincides with ground state energy. Symmetry breaking can occur. This is exactly the case when ρ is sufficiently large.

Theorem 0.1. *There exists $\rho_0 > 0$ such that $I_{\rho^2} < I_{\rho^2}^{rad}$ for all $\rho > \rho_0$.*

Remark 0.1. Our result does not clarify if minimizers for E restricted to B_ρ exist for large ρ . This is still an open problem.

Our argument is based on two ingredients. The first ingredient is the following scaling invariant inequality that holds only for radially symmetric function involving the kinetic term, the Coulomb term and L^p norms.

$$(0.2) \quad \|\varphi\|_{L^p(\mathbb{R}^3)} \leq C(p, s) \|\varphi\|_{\dot{H}^s(\mathbb{R}^3)}^{\frac{\theta}{2-\theta}} \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|\varphi(x)|^2 |\varphi(y)|^2}{|x-y|} dx dy \right)^{\frac{1-\theta}{4-2\theta}}$$

with $\theta = \frac{6-\frac{5}{2}p}{3-ps-p}$. Here the parameters s and p satisfy

$$p \in \left(\frac{16s+2}{6s+1}, \frac{6}{3-2s} \right] \text{ if } 1/2 < s < 3/2.$$

In our case we will use (0.2) in case $p = \frac{8}{3}$ and $s = 1$ to control from above the Slater correction term in terms of kinetic energy and Coulomb energy. In particular we obtain for radially symmetric functions that

$$(0.3) \quad \|\varphi\|_{L^{\frac{8}{3}}(\mathbb{R}^3)}^{\frac{8}{3}} \leq C \|\nabla \varphi\|_{L^2(\mathbb{R}^3)}^{\frac{4}{9}} \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|\varphi(x)|^2 |\varphi(y)|^2}{|x - y|} dx dy \right)^{\frac{5}{9}}$$

The importance of the previous inequality in the radial case is that it relates kinetic, Coulomb and Slater energy without information on the L^2 norm of the function.

The previous inequality will imply that $\inf_{\rho > 0} I_{\rho^2}^{rad} > -\infty$.

Remark 0.2. L^p lower bounds in terms of kinetic and Coulomb energy are not a novelty (see e.g. [2] for the general case). The first result concerning an improvement in radial case is due to Ruiz [14] (see also [13]) using a weighted Sobolev embedding for radially symmetric functions due to [16]. The scaling invariant radial bound (0.2) is proved in [3] using a pointwise inequality found by De Napoli [9] together with a lower bound for the Coulomb energy due to Ruiz [14].

The second ingredient is the asymptotic behaviour of the function

$$(0.4) \quad \rho \rightarrow \frac{I_{\rho^2}}{\rho^2}$$

for a translation invariant energy functional. We will show in Lemma 1.2 that the behaviour of $\rho \rightarrow \frac{I_{\rho^2}}{\rho^2}$ will imply that

$$I_{\rho^2} < I_{\mu^2} \text{ for all } 0 < \mu < \rho$$

$$\inf_{\rho > 0} I_{\rho^2} = -\infty$$

and hence our main result.

Remark 0.3. The importance of the function of $\rho \rightarrow \frac{I_{\rho^2}}{\rho^2}$ to show existence of minimizers is well established, see e.g. [6]. Indeed, the strict subadditivity inequality

$$I_{\rho^2} < I_{\mu^2} + I_{\rho^2 - \mu^2} \text{ for all } 0 < \mu < \rho$$

which implies the existence of minimizers follows immediately from the strict monotonicity of the function of $\rho \rightarrow \frac{I_{\rho^2}}{\rho^2}$. In the radial case, sufficient condition for the existence of minimizers is the strict monotonicity of $\rho \rightarrow I_{\rho^2}$. Here we show that the ratio between ground state energy and number of particles is crucial also for symmetry breaking.

Remark 0.4. We notice that our argument is general and symmetry breaking appears if we substitute the energy functional with

$$\tilde{E}(\psi) := \frac{1}{2} \|\nabla \psi\|_{L^2}^2 + \frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|\psi(x)|^2 |\psi(y)|^2}{|x - y|} dx dy - \frac{1}{p} \int_{\mathbb{R}^3} |\psi|^p dx$$

for $\frac{18}{7} < p < 3$. This fact follows again from the lower bounds for radially symmetric function (0.3). The exponent $p = \frac{18}{7}$ is the left endpoint exponent for which (0.3) holds.

Remark 0.5. We notice that $\lim_{\rho \rightarrow \infty} \frac{I_{\rho^2}^{rad}}{\rho^2} = 0$. Indeed, arguing as Theorem 1.2 in [14] or Theorem 2.2 in [3], there exist a family of functions $u_\rho \in B_\rho^{rad}$ with $\rho \rightarrow \infty$ with kinetic and Coulomb terms uniformly bounded and vanishing Slater term. This fact implies that $\sup_\rho I_{\rho^2}^{rad} < +\infty$.

Our last result concerns ground state energy in the radial case

Theorem 0.2. *The following alternative holds: either the function $\rho \rightarrow I_{\rho^2}^{rad}$ is monotone decreasing which implies the existence of minimizers for $E(u)$ on B_ρ^{rad} for all $\rho > 0$, or there exist a static radial solution to (0.1) belonging to $L^2(\mathbb{R}^3)$.*

Remark 0.6. The existence of static radial solution to (0.1) has been proved for the first time by Ruiz [14]. In [4] it has been shown that real solution to

$$(0.5) \quad -\Delta\psi + (|x|^{-1} * |\psi|^2)\psi - |\psi|^{p-2}\psi = 0$$

found by [11] belong always to $H^1(\mathbb{R}^3)$ if $3 \leq p < 6$. It remains open the case $p = \frac{8}{3}$ discussed here.

1. PROOF OF THEOREM 0.1.

Lemma 1.1. *Let $u \in B_{\rho^2}^{rad}$ then there exist $K_1 > 0$ not depending on ρ , such that*

$$E(u) > -K_1.$$

Proof. Let us call

$$x = \|\nabla u\|_{L^2(\mathbb{R}^3)}^2$$

$$y = \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|u(x)|^2 |u(y)|^2}{|x - y|} dx dy \right).$$

From inequality (0.3) one gets

$$E(u) > \frac{1}{2}x + \frac{1}{4}y - \frac{C}{p}x^{\frac{2}{9}}y^{\frac{5}{9}}$$

which immediately implies that $E(u)$ is bounded from below by a constant not depending on ρ .

In the general case, when we substitute the exponent $\frac{8}{3}$ for the Slater term with $\frac{18}{7} < p < 3$, we get

$$\tilde{E}(u) \geq \frac{1}{2}x + \frac{1}{4}y - \frac{C}{p}x^{\frac{5}{6}p-2}y^{1-\frac{p}{6}},$$

which again proves that the energy is bounded from below by a constant not depending on ρ .

□

We have the following general result for ground state energy of translation invariant energy functionals

Lemma 1.2. *Let the function $\rho \rightarrow I_{\rho^2}$ fulfills the following conditions*

$$(1.1) \quad -\infty < I_{\rho^2} < 0 \text{ for all } \rho > 0$$

the ground state energy is weakly subadditive, namely that

$$(1.2) \quad I_{\rho^2} \leq I_{\mu^2} + I_{\rho^2 - \mu^2} \text{ for all } 0 < \mu < \rho$$

$$(1.3) \quad \text{the function } \rho \rightarrow I_{\rho^2} \text{ is continuous}$$

$$(1.4) \quad \lim_{\rho \rightarrow 0} \frac{I_{\rho^2}}{\rho^2} = 0$$

then

$$I_{\rho^2} < I_{\mu^2} \text{ for all } 0 < \mu < \rho$$

$$\inf_{\rho > 0} I_{\rho^2} = -\infty$$

Proof. The strict monotonicity of the function $\rho \rightarrow I_{\rho^2}$ follows immediately from (1.1) and (1.2).

Let us argue now by contradiction assuming that there exist $K_2 > 0$ such that $I_{\rho^2} > -K_2$ for all $\rho > 0$. In this case we would have $\lim_{\rho \rightarrow \infty} \frac{I_{\rho^2}}{\rho^2} = 0$. By conditions (1.1), (1.3) and (1.4) the function $\rho \rightarrow \frac{I_{\rho^2}}{\rho^2}$ attains a global minimum, i.e. there exists ρ_0 such that

$$\frac{I_{\rho_0^2}}{\rho_0^2} \leq \frac{I_{\rho^2}}{\rho^2} \text{ for all } \rho > 0.$$

On the other hand, by the weak subadditivity condition (1.2),

$$\frac{I_{2\rho_0^2}}{2\rho_0^2} \leq \frac{I_{\rho_0^2} + I_{\rho_0^2}}{2\rho_0^2} = \frac{I_{\rho_0^2}}{\rho_0^2}$$

which implies that the function $\rho \rightarrow \frac{I_{\rho^2}}{\rho^2}$ attains a global minimum also at $\sqrt{2}\rho_0$. The same argument shows that

$$\frac{I_{k\rho_0^2}}{k\rho_0^2} = \frac{I_{\rho_0^2}}{\rho_0^2} \text{ for all } k \in \mathbb{N}$$

which implies that $\liminf_{\rho \rightarrow \infty} \frac{I_{\rho^2}}{\rho^2} < 0$ that contradicts the fact that $\lim_{\rho \rightarrow \infty} \frac{I_{\rho^2}}{\rho^2} = 0$. □

We show now that I_{ρ^2} fulfills the assumptions of Lemma 1.2, see also [6], [15].

Lemma 1.3. *The function $\rho \rightarrow I_{\rho^2}$ fulfills the following conditions*

$$-\infty < I_{\rho^2} < 0 \text{ for all } \rho > 0$$

$$I_{\rho^2} \leq I_{\mu^2} + I_{\rho^2 - \mu^2} \text{ for all } 0 < \mu < \rho$$

the function $\rho \rightarrow I_{\rho^2}$ is continuous

$$\lim_{\rho \rightarrow 0} \frac{I_{\rho^2}}{\rho^2} = 0$$

Proof. For generality we substitute the exponent $\frac{8}{3}$ for the Slater term with $\frac{18}{7} < p < 3$. The weak subadditive inequality is a general fact for translation invariant energy functionals, see e.g [15].

We define the following quantities:

$$A(u) := \int_{\mathbb{R}^3} |\nabla u|^2 dx, \quad B(u) := \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|u(x)|^2 |u(y)|^2}{|x - y|} dx dy \quad C(u) := - \int_{\mathbb{R}^3} |u|^p dx.$$

Negativity:

Let us consider, for $u \in B_1$, the rescaled function given by

$$u_{\theta, \beta}(x) = \theta^{1 - \frac{3}{2}\beta} u\left(\frac{x}{\theta^\beta}\right)$$

so that $\bar{\rho} := \|u_{\theta, \beta}\|_2 = \theta$. We easily find the following scaling laws:

$$A(u_{\theta, \beta}) = \theta^{2-2\beta} A(u),$$

$$B(u_{\theta, \beta}) = \theta^{4-\beta} B(u),$$

$$C(u_{\theta, \beta}) = \theta^{(1 - \frac{3}{2}\beta)p + 3\beta} C(u).$$

Notice that for $\beta = -2$ we get

$$E(u_{\theta, -2}) = \frac{\theta^6}{2} A(u) + \frac{\theta^6}{4} B(u) + \frac{\theta^{4p-6}}{p} C(u)$$

and we have $4p - 6 < 6$ since $p < 3$. Hence for $\theta \rightarrow 0$ we have $E(u_{\theta, -2}) \rightarrow 0^-$. This proves that there exists a small θ , and therefore a small $\bar{\rho}$, such that

$$I_{s^2} < 0 \quad \forall s \in (0, \bar{\rho}].$$

By weak subadditivity $I_{s^2} < 0$ for all s .

Continuity:

We first prove that if $\rho_n \rightarrow \rho$ then $\lim_{n \rightarrow \infty} E_{\rho_n^2} = I_{\rho^2}$. For every $n \in \mathbb{N}$, let $w_n \in B_{\rho_n}$ such that $E(w_n) < E_{\rho_n^2} + \frac{1}{n} < \frac{1}{n}$. Therefore, by using the interpolation and the Sobolev inequality, we get

$$\frac{1}{2} \|\nabla w_n\|_2^2 - C \rho_n^{\frac{6-p}{2}} \|\nabla w_n\|_2^{\frac{3(p-2)}{2}} \leq \frac{1}{2} \|\nabla w_n\|_2^2 - \frac{1}{p} \|w_n\|_p^p \leq E(w_n) < \frac{1}{n}.$$

Since $\frac{3(p-2)}{2} < 2$ and $\{\rho_n\}$ is bounded, we deduce that

$$\{w_n\} \text{ is bounded in } H^1(\mathbb{R}^3).$$

In particular $\{A(w_n)\}$ and $\{C(w_n)\}$ are bounded sequences, and also $\{B(w_n)\}$ since

$$\forall u \in H^1(\mathbb{R}^3) : B(u) \leq C\|u\|_{H^1(\mathbb{R}^3)}^4.$$

So we easily find

$$\begin{aligned} I_{\rho^2} &\leq E\left(\frac{\rho}{\rho_n}w_n\right) = \frac{1}{2}\left(\frac{\rho}{\rho_n}\right)^2 A(w_n) + \frac{1}{4}\left(\frac{\rho}{\rho_n}\right)^4 B(w_n) + \frac{1}{p}\left(\frac{\rho}{\rho_n}\right)^p C(w_n) \\ &= E(w_n) + o(1) < I_{\rho_n^2} + o(1). \end{aligned}$$

On the other hand, given a minimizing sequence $\{v_n\} \subset B_\rho$ for I_{ρ^2} , we have

$$I_{\rho_n^2} \leq E\left(\frac{\rho_n}{\rho}v_n\right) = E(v_n) + o(1) = I_{\rho^2} + o(1).$$

We get $\lim_{n \rightarrow \infty} I_{\rho_n^2} = I_{\rho^2}$.

Asymptotic behavior at zero:

In order to show that $\lim_{\rho \rightarrow 0} \frac{I_{\rho^2}}{\rho^2} = 0$, we notice that

$$\frac{G_{\rho^2}}{\rho^2} \leq \frac{I_{\rho^2}}{\rho^2} < 0$$

where G_{ρ^2} is defined by

$$(1.5) \quad G_{\rho^2} = \inf_{B_\rho} G(u)$$

where

$$G(u) = \frac{1}{2}\|\nabla u\|_{L^2(\mathbb{R}^3)}^2 - \frac{1}{p} \int_{\mathbb{R}^3} |u|^p dx.$$

It is well established that minimizers for G_{ρ^2} exist for all ρ (indeed it is not difficult to show that the function $\rho \rightarrow \frac{G_{\rho^2}}{\rho^2}$ is monotone decreasing). Let us call $\bar{u} \in B_{\rho^2}$ the minimizer for G_{ρ^2} , i.e the unique positive solution to

$$-\Delta u - \lambda_\rho u - |u|^{p-2}u = 0$$

where $\lambda_\rho < 0$ is the corresponding Lagrange multiplier. By Pohozaev identity it follows that

$$E(\bar{u}) = G_{\rho^2} = c\lambda_\rho \rho^2.$$

To conclude we shall prove that $\lim_{\rho \rightarrow 0} \lambda_\rho = 0$. Taken $\tilde{u}$ the unique positive solution to

$$-\Delta u + u - |u|^{p-2}u = 0$$

we have by scaling that $\bar{u} = (-\lambda_\rho)^{\frac{1}{p-2}} \tilde{u}(\sqrt{-\lambda_\rho}x)$ and hence

$$\rho^2 = \|\bar{u}\|_{L^2(\mathbb{R}^3)}^2 = (-\lambda_\rho)^{\frac{2}{p-2}-\frac{3}{2}} \|\tilde{u}\|_{L^2(\mathbb{R}^3)}^2$$

and noticing that $\frac{2}{p-2} - \frac{3}{2} > 0$ for $p < \frac{10}{3}$ we get the required estimate. $\square$

Proof of Theorem 0.1.

By Lemma 1.2 the ground state energy I_{ρ^2} is strictly decreasing as a function of ρ and $\inf_{\rho>0} I_{\rho^2} = -\infty$. On the other hand by Lemma 1.1, for radially symmetric functions $\inf_{\rho>0} I_{\rho^2}^{rad} > -\infty$ and therefore symmetry breaking occurs for sufficiently large ρ . $\square$

2. PROOF OF THEOREM 0.2.

The fact that the strict monotonicity of $\rho \rightarrow I_{\rho^2}^{rad}$ is sufficient for proving the existence of minimizers in the radial case follows immediately from the weak continuity of the Coulomb energy and of Slater energy. Indeed, taken a minimizing sequence $u_n \in B_{\rho^2}^{rad}$, $u_n \rightharpoonup u \neq 0$. Let us assume by contradiction that $u \in B_{\mu^2}^{rad}$ with $0 < \mu < \rho$, such that

$$I_{\mu^2}^{rad} + o(1) \leq E(u) + o(1) = E(u_n) = I_{\rho^2}^{rad} + o(1).$$

On the other hand, $I_{\mu^2}^{rad} > I_{\rho^2}^{rad}$ and therefore $\mu = \rho$ and u is a minimizer for E on $B_{\rho^2}^{rad}$.

It has been proved in [10] that $I_{\rho^2} = I_{\rho^2}^{rad}$ for sufficiently small ρ . On the other hand, by the weak subadditivity inequality it is clear that $I_{\rho^2} < I_{\mu^2}$ for $0 < \mu < \rho$. This fact proves that for sufficiently small ρ , $I_{\rho^2}^{rad} < I_{\mu^2}^{rad}$.

Let us define $c := \min_{[0,\rho]} I_{s^2}^{rad} < 0$ and

$$\rho_0 := \min \{s \in [0, \rho] \text{ s.t. } I_{s^2}^{rad} = c\}.$$

It is clear that $\rho_0 > 0$ and

$$(2.1) \quad \forall s \in [0, \rho_0] : I_{\rho_0^2}^{rad} < I_{s^2}^{rad}$$

namely, the function $[0, \rho_0] \ni s \mapsto I_{s^2} \in \mathbb{R}_-$ achieves the minimum only in $s = \rho_0$, by definition of ρ_0 . From this fact we deduce the existence of $\bar{u} \in B_{\rho_0}$ such that $E(\bar{u}) = I_{\rho_0^2}^{rad}$, and therefore $\bar{u}$ fulfills

$$(2.2) \quad -\Delta u + (|x|^{-1} * |u|^2)u - |u|^{\frac{2}{3}}u = 0$$

where λ is the associated Lagrange multiplier.

Now let us suppose that $\rho_0 < \rho$, which implies the existence of a sequence $\theta_n > 1$ with $\theta_n \rightarrow 1$ such that

$$E(\theta_n \bar{u}) \geq E(\bar{u}) \text{ with } \theta_n \rightarrow 1$$

and

$$E(\theta \bar{u}) \geq E(\bar{u}) \text{ with } 0 < \theta < 1.$$

This fact implies that

$$\frac{d}{d\theta}E(\theta\bar{u})|_{\theta=1} = 0,$$

and therefore the identity

$$(2.3) \quad \int_{\mathbb{R}^3} |\nabla \bar{u}|^2 dx + \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|\bar{u}(x)|^2 |\bar{u}(y)|^2}{|x-y|} dx dy - \int_{\mathbb{R}^3} |\bar{u}|^{\frac{8}{3}} dx = 0.$$

From (2.3) we notice that the associated Lagrange multiplier in (2.2) is $\lambda = 0$, and hence $\bar{u}$ solves

$$-\Delta u + (|x|^{-1} * |u|^2)u - |u|^{\frac{2}{3}}u = 0,$$

proving that $\bar{u}$ is a static solution to (0.1) with $\bar{u} \in B_{\rho_0^2}^{rad}$.

REFERENCES

- [1] C. Bardos, F. Golse, A. D. Gottlieb, N. Mauser, Mean field dynamics of fermions and the time-dependent Hartree-Fock equation, *J. Math. Pures Appl.* (9) 82 (2003), no. 6, 665-683.
- [2] J. Bellazzini, R.L. Frank, N. Visciglia, Maximizers for Gagliardo-Nirenberg inequalities and related non-local problems, *Math. Annalen* (2014), no. 3-4, 653-673.
- [3] J. Bellazzini, M. Ghimenti, T. Ozawa, *Sharp lower bounds for Coulomb energy*, *Math. Res. Lett.* (in press) arXiv:1410.0598 (2015)
- [4] J. Bellazzini, L. Jeanjean, T. Luo, Existence and instability of standing waves with prescribed norm for a class of Schrödinger-Poisson equations, *Proc. London Math. Soc.* (3) 107,(2013), 303-339.
- [5] J. Bellazzini, G. Siciliano, Stable standing waves for a class of nonlinear Schrödinger-Poisson equations, *Z. Angew. Math. Phys.*, 62 (2011), no. 2, 267-280.
- [6] J. Bellazzini, G. Siciliano, Scaling properties of functionals and existence of constrained minimizers, *J. Funct. Analysis*, 261 (2011), no. 9, 2486-2507.
- [7] O. Bokanowski, J.L. Lopez, O. Sanchez, J. Soler, Long time behaviour to the Schrödinger-Poisson- X^α systems. *Mathematical physics of quantum mechanics*, 217-232, *Lecture Notes in Phys.*, 690, (2006)
- [8] I. Catto, J. Dolbeault, O. Sanchez, J. Soler, Existence of steady states for the Maxwell-Schrodinger-Poisson system: exploring the applicability of the concentration-compactness principle, *Math. Models Methods Appl. Sci.* 23 (2013), no. 10, 1915-1938.
- [9] P.L. De Nápoli, Symmetry breaking for an elliptic equation involving the Fractional Laplacian, arXiv:1409.7421
- [10] V. Georgiev, F. Prinari, N. Visciglia, On the radiality of constrained minimizers to the Schrödinger-Poisson-Slater energy, *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2012), no. 3, 369-376.
- [11] I. Ianni, D. Ruiz, Ground and bound states for a static Schrödinger-Poisson-Slater problem. *Commun. Contemp. Math.* 14 (2012), no. 1, 1250003
- [12] N. J. Mauser, The Schrödinger-Poisson- X^α equation, *Appl. Math. Lett.* 14 (2001), no. 6, 759-763.
- [13] D. Ruiz, The Schrödinger-Poisson equation under the effect of a nonlinear local term, *J. Funct. Anal.* 237 (2006) 655-674.
- [14] D. Ruiz, On the Schrödinger-Poisson-Slater system: behavior of minimizers, radial and nonradial cases, *Arch. Ration. Mech. Anal.* 198 (2010), no. 1, 349-368

- [15] O. Sanchez, J. Soler, Long-time dynamics of the Schrödinger-Poisson-Slater system, J. Statist. Phys., 114 (2004), no. 1-2, 179-204.
- [16] J. Su, Z.-Q. Wang, M. Willem, Weighted Sobolev embedding with unbounded and decaying radial potentials, J. Differential Equations 238 (2007), 201-219.

UNIVERSITÀ DI SASSARI, VIA PIANDANNA 4, 07100 SASSARI, ITALY
E-mail address: jbellazzini@uniss.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO B. PONTECORVO 5, 56100
PISA, ITALY
E-mail address: ghimenti@mail.dm.unipi.it